

CT Children's CLASP Guideline

Post-COVID Conditions and COVID-19 Long Haul Management

INTRODUCTION

As of September 2021, the COVID-19 pandemic has infected over 5 million children (of over 33 million infections worldwide), representing over 15% of all COVID-19 cases. This number continues to rise with the delta variant of SARS-CoV-2. Although pediatric hospitalizations and deaths remain relatively low, children may be left with long-lasting, post-acute sequelae from their infection, even if the child was initially asymptomatic. "Post-COVID Conditions" is a term to describe the wide range of physical and mental health consequences that are present for ≥ 4 weeks after an initial COVID-19 infection, even in children who were initially asymptomatic (CDC, 2021). Post-COVID Conditions are thought to affect 5% to 80% of individuals who have had COVID-19 infection of any severity, but the data is limited on its impact on the pediatric population. Because of the uncertainty surrounding Post-COVID Conditions, including the wide range of presentations, difficulty in assessing potential alternative etiologies, and lack of standardized guidance, a thorough, systematic and multi-disciplinary approach to evaluation and management is warranted.

Important note: It is important to distinguish between the care outlined below for Post-COVID Conditions and that of **Multisystem Inflammatory Syndrome in Children (MIS-C)**, which can appear weeks after a child's initial infection as well. MIS-C can be a medical emergency and the management should not follow this referral guideline. The CDC defines MIS-C as an individual < 21 years of age with fever, laboratory evidence of inflammation, and evidence of clinically severe illness requiring hospitalization, with multisystem (≥ 2) organ involvement (cardiac, renal, respiratory, hematologic, gastrointestinal, dermatologic, or neurological) AND no alternative plausible diagnosis; AND positive for current/recent SARS-CoV-2 infection (or exposure to suspected/confirmed COVID-19 case within the 4 weeks prior to the onset of symptoms).

INITIAL EVALUATION

INITIAL EVALUATION:

PCPs should be mindful of the presence of post-COVID conditions after any documented SARS-CoV-2 infection, even if the child is initially asymptomatic. New or persisting symptoms are present for ≥ 4 weeks from acute infection (i.e., ≥ 4 weeks from symptom onset and/or first positive test). Note that patients who present with possible MIS-C should be excluded from this guideline.

RELEVANT CLINICAL HISTORY/REVIEW OF SYSTEMS:

- **Past Medical History:**

- Confirmation of SARS-CoV-2 infection and date of onset
- Symptom severity (if any) during acute infection
- History of thrombosis
- Other physical or mental underlying disease states present that may contribute to presentation

- **Family History:**

- Note if any immediate family with inherited thrombophilia (i.e., Factor V Leiden, Prothrombin Gene Mutation, Protein S or Protein C deficiency, Anti-thrombin III deficiency)

- **Symptoms Associated with Post-COVID Conditions (most common):**

****Symptoms present for ≥ 4 weeks from acute infection (i.e., ≥ 4 weeks from symptom onset and/or first positive test)****

- **Overall Wellness:**

- Post-exertional malaise and/or poor endurance (worsening of symptoms following minor physical/mental exertion, worsening 12-48 hours after activity and lasting for days or even weeks)

- **Cognitive/Mental:**
 - Insomnia/difficulty sleeping or hypersomnia
 - Fatigue (compared to before COVID-19 infection)
 - Depression, anxiety, or mood/behavioral changes compared to before COVID-19 infection
 - Brain fog, cognitive impairment
 - Complete [Appendix A: Pediatric Cognitive Function – Short Form](#)
- **Head:**
 - Mild-moderate, intermittent headache
 - Seizures
- **ENT:**
 - Anosmia (loss of smell)
 - Change or loss of taste
 - Persistent nasal congestion
- **Lungs:**
 - Intermittent dyspnea
 - Persistent cough
 - Exercise intolerance
 - Chest tightness
- **Heart:**
 - Palpitations and/or tachycardia
 - Lightheadedness
 - Chest pain
- **GI:**
 - Abdominal pain
 - Diarrhea
 - Constipation
 - Nausea/vomiting or inability to tolerate intake
- **GU:**
 - Difficulty or painful urination
- **MS:**
 - Pain
 - Arthralgia
 - Myalgia
 - Paresthesia, neuropathy
 - Muscle weakness
 - Abnormal movements
- **Skin:**
 - Rash or abnormal discoloration
- **Endocrine:**
 - Menstrual cycle irregularities
 - Significant weight loss or weight gain
- **Infectious:**
 - Fever *without concern for MIS-C*
- **Heme:**
 - Signs of thrombosis or anemia
- **Other:**
 - Any new symptoms that have persisted for more than 4 weeks after a confirmed SARS-CoV-2 infection

- **Timing of Onset:**
 - Confirm timing of initial infection
 - Note persisting or new symptoms that are present ≥ 4 weeks after initial acute infection (i.e., ≥ 4 weeks from symptom onset and/or first positive test) that are not easily explained by an alternative etiology
- FULL PHYSICAL EXAM:**
- *A full physical examination should be completed by the PCP and should note the following components:**
- Vitals: HR, RR, O2 sat, BP, temperature
 - Anthropometrics: Weight, height, BMI
 - Skin: Full skin examination; include any swelling, pigmentation, varicosities, ulcers
 - HEENT: standard examination; include any oral ulcerations
 - Cardiac: include heart sounds, perfusion, liver edge, edema
 - Lung: include SpO2, location/presence of abnormalities
 - GI: include presence of hepatosplenomegaly, distention, abnormal bowel sounds
 - GU: standard examination, if needed
 - MS: include joint examination, ROM, pain, edema
 - Neuro: include a gross neurological examination
 - Heme: assess for any indication of thrombosis, DVT, PE, or stroke
- **RED FLAGS:**
 - Any abnormalities on examination should be further investigated and referred, as appropriate
 - **URGENT/EMERGENT RED FLAGS:**
 - Any symptoms or signs of clinical instability, including, but not limited to:
 - severe chest pain, severe shortness of breath, severe persistent headache, severe abdominal pain, persistent fever with concern for MIS-C, signs of thrombosis/DVT/PE/stroke
 - active suicidality or mental health instability placing child at immediate risk of harm to self or others

INITIAL PCP MANAGEMENT

- Initial Management by PCP:**
- Establish if the patient needs urgent/emergent referral (see URGENT/EMERGENT RED FLAGS above):
 - Refer patient to emergency care, as appropriate
 - Establish if acute intervention is needed (see RED FLAGS above):
 - Standard clinical management should be initiated by the PCP on a case-by-case basis of patient’s presenting signs and symptoms
 - Establish if above signs and symptoms could be due to re-infection with SARS-CoV-2:
 - Although rare, reinfection within 90 days of initial infection could occur
 - Consider symptoms, risk factors/exposures, and discussing case with CT Children’s Infectious Diseases providers

WHEN TO REFER TO LONG HAUL CLINIC

- All patients with a concern for post-COVID conditions can be referred to CT Children’s “Long Haul” Clinic in Infectious Diseases. All referrals are considered routine and will be scheduled in 2-3 weeks.
- **Routine referral to Long Haul Clinic if all of the following criteria met:**
 - Documented SARS-CoV-2 infection ≥ 4 weeks prior
 - New or persisting symptoms listed above are present for ≥ 4 weeks from acute SARS-CoV-2 infection (i.e., ≥ 4 weeks from symptom onset and/or positive test result)
 - Abnormal/positive **Pediatric Cognitive Screen: Appendix A**

	<ul style="list-style-type: none"> • Defer referral if any of the following criteria are present: <ul style="list-style-type: none"> ○ Absence of acute SARS-CoV-2 infection ○ New or persisting symptoms have been present for <4 weeks (i.e., <4 weeks from symptom onset and/or first positive test result) ○ New or persisting symptoms are easily attributable to alternative, underlying disease process • Emergency management if URGENT/EMERGENT RED FLAGS are present. Routine referrals are not clinically appropriate in these instances.
<p>HOW TO REFER</p>	<p>Routine Referral to Department of Infectious Diseases and Immunology via CT Children’s One Call Access Center Phone: 833.733.7669 Fax: 833.226.2329</p> <p>Information to be included with the referral:</p> <ul style="list-style-type: none"> ▪ Notes from recent visits that document specific Post-COVID Conditions that are present ▪ Growth charts (height, weight, BMI) ▪ Copy of documented laboratory report of SARS-CoV-2 infection (include date of test) ▪ Copy of Pediatric Cognitive Function – Short Form (Appendix A) ▪ Copies of other relevant laboratory studies and imaging studies
<p>WHAT TO EXPECT – PCPs</p>	<p>What to expect from CT Children’s Visit:</p> <ul style="list-style-type: none"> ▪ Initial evaluation by Long Haul Clinic in Infectious Diseases within 2-3 weeks. This may be an in-person or telemedicine visit. ▪ Comprehensive history, review of systems and physical exam ▪ Confirmation of Long Haul diagnosis by Infectious Diseases provider <ul style="list-style-type: none"> ○ Baseline laboratory studies ○ Additional laboratory and imaging studies, as indicated ○ Referral to appropriate subspecialists to assess for other organic causes, as appropriate ○ If appropriate, patient will be referred to CT Children’s Pain Team to assist with a multi-disciplinary approach for symptomatic management of certain symptoms such as fatigue, brain fog, etc. ▪ If Long Haul diagnosis is unlikely to be the etiology of current presentation, patient will be transitioned back to the PCP for further management.
<p>WHAT TO EXPECT – Patients/Families/Caregivers</p>	<p>PCPs to review with patients/families/caregivers:</p> <ul style="list-style-type: none"> ▪ Initial evaluation by Long Haul Clinic in Infectious Diseases within 2-3 weeks, to confirm Post-COVID Condition diagnosis. There is no specific cure for Post-COVID Conditions and much is still unknown. Your child will be managed on a case-by-case basis, and resolution of symptoms may take some time. Symptoms may initially be managed by your primary care provider. ▪ Your initial visit with the Long Haul Clinic may be in-person or through a telemedicine visit. ▪ The provider may order further lab tests and studies, as appropriate. ▪ If appropriate, your symptoms may be managed by a multi-disciplinary group of specialists, with further specialty follow up to assess for any other causes of your symptoms. It may take 2-3 weeks to be seen by other specialists after your initial visit with Infectious Diseases. ▪ Your specialists will be communicating closely with your primary care provider.

Appendix A: Pediatric Cognitive Function – Short Form

Screening is positive if: score <25 or ≥2 items are “quite/very much”.

If patient has baseline ADHD, a significant change from baseline (prior to COVID-19 infection) should prompt a referral.

Neuro-QOL Item Bank v2.0 –Pediatric Cognitive Function – Short Form

Pediatric Cognitive Function – Short Form

Please respond to each question or statement by marking one box per row.

		Not at all	A little bit	Somewhat	Quite a bit	Very much
NCCOGped03	I forget schoolwork that I need to do.....	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1
NCCOGped05	I sometimes forget what I was going to say.....	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1
NCCOGped08	I react slower than most people my age when I play games.	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1
NCCOGped15	I forget things easily.....	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1
NCCOGped17	I have trouble remembering to do things (e.g., school projects).....	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1
NCCOGped18	It is hard for me to concentrate in school.....	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1
NCCOGped19	I have trouble paying attention to the teacher.....	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1
NCCOGped20	I have to work really hard to pay attention or I will make a mistake.....	<input type="checkbox"/> 5	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1

©2008-2013 David Cella and the PROMIS Health Organization on behalf of the National Institute for Neurological Disorders and Stroke (NINDS). Used with permission.

Key References:

- [Children and COVID-19: State-Level Data Report \(aap.org\)](#)
- [Key Points | Evaluating and Caring for Patients with Post-COVID Conditions | CDC](#)
- [Tracking - Johns Hopkins Coronavirus Resource Center \(jhu.edu\)](#)

Other References:

- Brackel, C.L.H., Lap, C.R., Buddingh, E.P., et al. Pediatric long-COVID: an overlooked phenomenon?. *Pediatr Pulmonol*. Published online June 8, 2021. doi:[10.1002/ppul.25521](#).
- Buonsenso, D., Munblit, D., De Rose, C., et al. Preliminary evidence on long COVID in children. *Acta Paediatr*. 2021;110(7):2208-2211. doi:[10.1111/apa.15870PubMed](#).
- Ludvigsson, J.F. Case report and systematic review suggest that children may experience similar long-term effects to adults after clinical COVID-19. *Acta Paediatr*. 2021;110(3):914-921. doi:[10.1111/apa.15673PubMed](#).
- Say, D., Crawford, N., McNab, S., Wurzel, D., Steer, A., Tosif, S. Post-acute COVID-19 outcomes in children with mild and asymptomatic disease. *Lancet Child Adolesc Health*. 2021;5(6):e22-e23. doi:[10.1016/S2352-4642\(21\)00124-3PubMed](#).