INTRODUCTION

Heart murmur is a common issue in pediatric patients, but the presence of a murmur does not always indicate the presence of heart disease. Pathologic murmurs include those louder than grade 3, all diastolic murmurs, unusually harsh systolic murmurs, and murmurs associated with clicks or abnormal splitting of S2. A murmur that increases in intensity with standing could indicate hypertrophic cardiomyopathy. Murmurs associated with dysmorphic features should be referred as there is a higher risk of congenital heart disease in syndromic patients. A number of innocent murmurs exist which are described below, and which, in the asymptomatic patient, do not require further evaluation by a pediatric cardiologist. The most common innocent murmurs are described in the table below:

	Type of Innocent Murmur	Typical Age Range	Key Features	Outcome(s)
	PERIPHERAL PULMONIC STENOSIS	Infants 1-3 months	High-pitched, systolic, heard at left upper sternal area and on back or in axillae	Unless infant is dysmorphic, or symptomatic, watchful waiting is appropriate – will go away within a few months
	STILL'S VIBRATORY MURMUR	Pre-schoolers and school age children; Occasionally heard in newborns	Low-pitched "twang"; best heard with the bell at lower-left sternal border when patient supine. Gets louder with increased cardiac output (fever, anemia, etc.)	Can come and go depending on state of the child, can enhance the murmur with brief exercise. Usually resolves by adolescence
	VENOUS HUM	Pre-school and young school-age	Continuous "windy" noise heard near the clavicles in a sitting patient Goes away with head turned to the side, with gentle pressure over the jugular vein, or when patient is supine	Disappears before adolescence
	PULMONARY FLOW MURMUR	Older school-age children and teens	Soft systolic "flow" murmur best heard at left upper sternal border, may get louder with anemia, exercise, fever	May be difficult to distinguish from an ASD or mild pulmonary valve stenosis. Refer if there is wide, fixed split S2, if a click is heard, or if murmur is harsh sounding
NITIAL VALUATION ND IANAGEMENT	 INITIAL EVALUATION: Obtain patient and family history Evaluate for RED FLAGS: Chest pain (especially with exercise), syncope or respiratory symptoms. In infants, look for fatigue and diaphoresis while feeding, cyanosis, poor weight gain. Obtain targeted physical exam Check femoral pulses and obtain blood pressure 			

- of the aorta
- Categorize murmur using above chart



	INITIAL MANAGEMENT:			
	 If applicable, explain diagnosis of innocent murmur to patient/family 			
	 Provide reassurance 			
	 Re-evaluate at next well child visit 			
	 If murmur does not easily fall into one of the innocent categories above, consider referral to Cardiology (see below) 			
WHEN	URGENT REFERRAL:			
TO REFER	 Asymptomatic babies less than 2 months of age will be seen by Cardiology in a timely manner. 			
	 If any symptoms are present in a newborn, contact Cardiologists immediately via One Call to determine the most appropriate triage. The risk of finding a critical ductal dependent lesion is greatest in the first 2 weeks. 			
	 Any patient with RED FLAGS (chest pain, palpitations, exercise intolerance, syncope). If symptoms are particularly concerning and a more emergent referral needed, please contact a cardiologist directly via One Call. 			
	ROUTINE REFERRAL:			
	 Any asymptomatic child >2 months whose murmur does not fit one of the "innocent" categories above 			
	 Any syndromic/dysmorphic patient (Down's, Marfan's, Fetal Alcohol, DiGeorge, etc.) 			
	 Any child for which parent anxiety persists despite reassurance 			
	 If primary care provider is unsure that murmur is innocent 			
HOW	Referral to Cardiology via CT Children's One Call Access Center			
TO REFER	Phone: 833.733.7669 Fax: 833.226.2329			
	For more information on how to place referrals to Connecticut Children's, click here.			
	Information to be included with the referral:			
	 Relevant findings on history and physical exam, including timing of onset of murmur 			
	 Growth chart, labs if any were done 			
	 Please DO NOT obtain echo prior to consultation. Studies done in an adult echo lab are not designed to look for congenital defects, and will often miss things like patent ductus, 			
	coarctation, or anomalous pulmonary veins. Cardiologist will determine if echo is warranted at time of initial consult.			
WHAT TO	What to expect from CT Children's Visit:			
EXPECT	 Meet with cardiologist to review patient and family history 			
	Physical exam			
	 EKG – done for all routine visits 			
	 Please tell families they will receive a cardiology consultation, and if the cardiologist 			
	determines that an echocardiogram is warranted, it will be arranged in a timely manner			

