



KNEE OSTEOCHONDritis DISSECANS (OCD) REPAIR POST-OPERATIVE INSTRUCTIONS AND HOME EXERCISE

The following information is to help make your recovery as smooth and rapid as possible.

1. Keep your knee elevated as much as possible for the next few days. Use crutches at all times.
2. Keep your dressing on for three days. Do not get it wet. You may shower by wrapping plastic wrap over your dressing. After three days, the bandages may be removed and the wounds covered with clean dressings. Keep the wounds dry until your first visit after surgery.
3. Post-operative bleeding is not unusual. Reinforce the dressing as needed. If you have concerns about the amount of bleeding, please call.
4. Use your cold therapy unit as directed until you are seen in the office.
5. Remember that the surgery will cause your thigh muscles to be weak, so take your time and be safe. You may rest your foot on the ground for balance only.
6. You may begin the exercises on page two. There is no limit to the amount of these exercises you may do. Stop any exercise that causes sharp pain.
7. Post-operative pain is common but should be controlled by the prescriptions given to you.
8. You will be seen for follow-up within a week. Please call our office if you have any questions or concerns.



KNEE OSTEOCHONDritis DISSECANS (OCD) REPAIR POST-OPERATIVE INSTRUCTIONS AND HOME EXERCISE



Heel-Leg Prop

Elevate your leg by placing a cushion underneath your lower leg or heel as shown. Do this 4-8 times per day for 15-20 minutes. You may also sleep with your leg elevated.



Quad Set

Squeeze thigh muscles tightly, as if pressing the back of your knee into the surface you are lying on. Hold contraction for 10 seconds, repeat 30 times. Do this 4-6 times per day.



Straight Leg Raise - Flexion

Squeeze thigh muscles tightly, then lift leg off the surface as shown in the picture. Raise and lower your leg slowly. Repeat 15-30 times. Do this 4-6 times per day.



Straight Leg Raise - Adduction

Squeeze thigh muscles tightly, then lift leg off the surface as shown in the picture. Raise and lower your leg slowly. Repeat 15-30 times. Do this 4-6 times per day.



Straight Leg Raise – Abduction

Squeeze thigh muscles tightly, then lift leg off the surface as shown in the picture. Raise and lower your leg slowly. Repeat 15-30 times. Do this 4-6 times per day.



Straight Leg Raise – Extension

Squeeze thigh muscles tightly, then lift leg off the surface as shown in the picture. Raise and lower your leg slowly. Repeat 15-30 times. Do this 4-6 times per day.



KNEE OSTEOCHONDritis DISSECANS (OCD) REPAIR POST-OPERATIVE INSTRUCTIONS AND HOME EXERCISE



Assisted Knee Flexion

While seated, allow your knee to bend a comfortable amount. Use your other foot to control bending and straightening your knee. Stop if this becomes uncomfortable. This may be done several times a day.



Calf Stretch

Using a towel or a belt, gently stretch your calf muscles as pictured. Keep your knee straight. Hold the stretch for 20-30 seconds. Do this 4-6 times per day.